

Hyperon polarization in ν_μ charged current interaction at the NOMAD energies

Liang Zuo-tang^{1,3}, and Liu Chun-xiu^{1,2}

¹*Department of Physics, Shandong University, Jinan, Shandong 250100, China*

²*Institute of High Energy Physics, The Chinese Academy of Sciences, Beijing 100039, China*

³*The Abdus Salam International Center for Theoretical Physics, 34100 Trieste, Italy*

Abstract

We show that, in $\nu_\mu N \rightarrow \mu^- \Lambda X$ at the NOMAD energies, it is impossible to separate the products of the fragmentation of the struck quark from those of the nucleon remnant. The latter has a large contribution even in the current fragmentation region and has to be taken into account in calculating Λ polarization using different pictures. Based on this, we make a rough estimation for the longitudinal Λ polarization in $\nu_\mu N \rightarrow \mu^- \Lambda X$ at the NOMAD energies. A comparison with the data is given and predictions for $\nu_\mu N \rightarrow \mu^- \Sigma^+ X$ are presented.

Longitudinal hyperon polarizations in lepton induced reactions have attracted much attention recently (See, e.g., [1-15] and the references given there) since they provide an important tool to study the spin transfer in high energy fragmentation processes. Spin transfer in fragmentation is defined as the probability for the polarization of the fragmenting quark to be transferred to the produced hadron. It is one of the important issues in the spin effects in the fragmentation processes. The study is of particular interest because it is still unclear whether the SU(6) or the DIS picture is suitable for the description of this problem. Hyperon polarization in lepton induced reactions is an ideal place to study this problem since here the polarization of the fragmenting quark can easily be calculated and the hyperon polarization can easily be measured by measuring the angular distribution of its decay products. At sufficiently high energies the hadrons produced in the current fragmentation region of the deeply inelastic lepton-nucleon scattering can be regarded as being completely from the struck quark fragmentation. Therefore it can be used to study the spin transfer from the fragmenting quark to the produced hadron. Theoretical calculations using different models have been made and predictions have been given^{4,9-11}.

It is encouraging to see that measurements of Λ polarization in deeply inelastic scattering have been carried out by HERMES at HERA¹³ and NOMAD at CERN¹⁴. They respectively measured the Λ polarization in $e^+p \rightarrow e^+\Lambda X$ at $E_e = 27.6$ GeV and in $\nu_\mu N \rightarrow \mu^-\Lambda X$ at, on the average, $E_\nu = 44$ GeV. In particular, NOMAD has measured it with rather high accuracy and has obtained the P_Λ 's for both target and current fragmentation regions, hence also its x_F dependence. [Here, $x_F \equiv 2p_{\parallel}/W$, $p_{\parallel}$ and W are respectively the momentum component of the produced hyperon parallel to the virtual boson direction and the total energy of the produced hadronic system in the center of mass (c.m.) frame of the system]. However, if we compare the NOMAD data¹⁴ with the theoretical predictions in [4,9-11], we see the following distinct difference: While all the theoretical results go to zero when x_F goes to zero, the data show that $|P_\Lambda|$ rises monotonically when x_F decreases from positive to negative. It does not go to zero when x_F goes to zero. What does this tell us? Does it imply that none of the pictures in [4,9-11] for spin transfer in high-energy fragmentation processes is applicable in

$$\nu_\mu N \rightarrow \mu^- \Lambda X?$$

In this note, we show that the answer to the question should be “No!” This is because, in all of the above-mentioned calculations^{4,9–11}, only the struck quark fragmentation has been taken into account and the influence from the fragmentation of the remnant of the scattered nucleon was neglected. This is a good approximation only at high energies. In contrast, in the NOMAD or HERMES energy region, the influence of the fragmentation of the nucleon remnant is usually very large. We have to take them into account in particular in the region near $x_F = 0$. We will show our results from a Monte-Carlo calculation using the event generator LEPTO¹⁶ and make a rough estimation of the Λ polarization by taking the fragmentation of the nucleon remnant into account using a valence quark model.

We recall that, in deeply inelastic lepton-nucleon scattering, we are usually envisaged with the following picture: During the collision, a quark (or an anti-quark) was struck by the exchanged virtual boson. Viewed in the c.m. frame of the hadronic system, the struck quark (anti-quark) flies in the opposite direction as the nucleon remnant. The system, which consists of the struck quark and the nucleon remnant, then fragments into the produced hadrons. At sufficiently high W , the hadronization products which contain the struck quark and those which contain one or more of the valence quarks of the nucleon remnant are usually the leading particles in the two opposite directions. Their x_F ’s are usually large in magnitude and opposite in sign so that they are well separated from each other. When we study the hyperon polarization in the large and positive x_F region, i.e. in the current fragmentation region, we need only to take the fragmentation of the struck quark into account but the influence of the nucleon remnant can be neglected. However, in e.g. the NOMAD experiments, where $\nu_\mu N \rightarrow \mu^- \Lambda X$ is studied, the incident energies of ν_μ lies between 10 and 50 GeV, and W is only several GeV. In this case, the number of the produced hadrons is usually very small. The $|x_F|$ for the hadrons which contain the struck quark or one or more of the valence quarks of the nucleon remnant can also be very small. It is even possible that they move in the opposite direction as the quark did before the fragmentation. As a consequence, it is impossible to separate the fragmentation products of

the struck quark and those of the nucleon remnant from each other, even at relatively large x_F . In particular, in the region of $x_F \sim 0$, the contributions from the nucleon remnant can be very important.

To study this effect explicitly, we now take $\nu_\mu p \rightarrow \mu^- \Lambda X$ as an example. At the relatively low energies, we consider only the valence quarks of proton thus are envisaged with the following picture: In the collision, a d -quark in proton is knocked out and converted to a u -quark, which flies in the opposite direction as the remaining (uu) -diquark. This u -(uu) system then fragments into hadrons. There are following possibilities to produce a Λ which contains one of the three u -quarks:

- (1) directly produced and contains the struck u ;
- (2) directly produced and contains one of the u from the (uu) -diquark;
- (3) decay product of a directly produced hyperon resonance that contains the struck u ;
- (4) decay product of a directly produced hyperon resonance that contains the two u 's or one of them from the (uu) -diquark.

In the cases (1) and (3), the vacuum excitation of a diquark-anti-diquark pair is needed. There should be at least one more baryon and an anti-baryon produced, i.e., there should be at least two baryons and one anti-baryon produced in the reaction. In the cases (2) and (4), there is no need for such vacuum excitation and there can be only one baryon produced. Since the total c.m. energy of the u -(uu)-system W is only several GeV, the probabilities for the cases (1) and (3) should be much more suppressed compared with those for the cases (2) and (4). Also because the energy of any one of the u 's in the (uu) -diquark is quite low, the Λ produced in case (2) and (4) may not necessarily fly in the same direction as the original u -quark. It can also be in the opposite direction. In particular in the $x_F \sim 0$ region, the contributions from the cases (2) and (4) can dominate.

Since whether the cases (2) and (4) dominate and, if yes, how strongly they dominate the Λ production in $\nu_\mu p \rightarrow \mu^- \Lambda X$, are questions independent of the spin transfer in fragmenta-

tion process, we can study them using a hadronization model which give a good description of the properties of the hadrons produced in unpolarized reactions. We thus use the event generator LEPTO¹⁶ based on Lund model¹⁷ to make the numerical estimations. We generated about 10^7 $\nu_\mu p \rightarrow \mu^- \Lambda X$ events, analyzed the origins of the produced Λ 's and obtained the results shown in Fig.1. From the figure, we see clearly that the contributions from the cases (2) and (4) are important. They are much higher than those from the cases (1) and (3) in the region near $x_F = 0$. Hence, to calculate the polarization of Λ in $\nu_\mu p \rightarrow \mu^- \Lambda X$ at such energies, in particular for $x_F \sim 0$, we have to take the fragmentation of the nucleon remnant into account.

Having seen that the contributions from the nucleon remnant are indeed very important in the NOMAD energy region, we now calculate the Λ polarization in $\nu_\mu N \rightarrow \mu^- \Lambda X$ by taking them into account. Apparently, the detailed results should depend very much on the polarizations of the quarks in the nucleon remnant, which are unclear yet. It is thus impossible to make a detailed calculation at present. However, since the energies are relatively low, the characteristic features of the results should be mainly determined by the valence quarks. We therefore make a rough estimation using a valence quark model to calculate the polarizations of the quarks in the nucleon remnant in the following.

Since we take only the valence quarks in the nucleon into account, the leading order hard subprocess in $\nu_\mu N \rightarrow \mu^- HX$ can only be $\nu_\mu d \rightarrow \mu^- u$. According to the standard model for electro-weak interaction, the polarization of the outgoing struck quark is $P_u^{(stru)} = -1$. We neglect the mass of the quarks, hence the helicity of the quark is conserved in the process $\nu_\mu d \rightarrow \mu^- u$. So the d -quark was also in the helicity “−” state before the scattering. For the reaction $\nu_\mu p \rightarrow \mu^- HX$, the nucleon remnant is a (uu) -diquark. If the proton was in the helicity “+” state, the remaining (uu) -diquark has to be in $(uu)_{1,-1}$ state. (Here we use the subscripts to denote the spin and its component along the moving direction of the outgoing struck quark in the c.m. frame of the hadronic system. The minus sign is because the moving direction of the outgoing struck quark is opposite to that of the diquark.) The relative probability, obtained from the $SU(6)$ wave function, is $2/3$. If the

proton is in the helicity “−” state, the remaining (uu) -diquark has to be in $(uu)_{1,0}$ state, and the relative probability is $1/3$. Hence, the u -quark in the nucleon remnant is polarized and the polarization is $P_u^{(di,p)} = -2/3$. For the reaction $\nu_\mu n \rightarrow \mu^- HX$, the nucleon remnant is a (ud) -diquark. Similarly, we obtain that this (ud) -diquark can be in the $(ud)_{1,-1}$, $(ud)_{1,0}$ or $(ud)_{0,0}$ state, and the relative probability is $1/6$, $1/12$ or $3/4$ respectively. Hence, the polarization of the u - or d -quark in the neutron remnant is $P_u^{(di,n)} = P_d^{(di,n)} = -2/3$ if it is from the spin-1 (ud) -diquark or $P_u^{(di,n)} = P_d^{(di,n)} = 0$ if it is from the spin-0 (ud) -diquark.

All these polarizations can be transferred to the hyperons produced in the hadronization. We calculate the polarizations for the hyperons of the different origins in the following ways: For those which are directly produced and contain the struck quark or one of the quarks in the nucleon remnant, we calculate in the same way as we did in [3,8,10,12] for $q \rightarrow HX$ using two different pictures, i.e. the SU(6) and the DIS pictures. For those which are directly produced and contain the diquark, we calculate using the SU(6) wave-function. The polarization of the Λ in the final state in $\nu_\mu p \rightarrow \mu^- \Lambda X$ is given by,

$$P_\Lambda = [P_u^{(stru)} t_{\Lambda,u}^F \langle n_\Lambda^{(1)} \rangle + P_u^{(di,p)} t_{\Lambda,u}^F \langle n_\Lambda^{(2)} \rangle + \sum_j t_{\Lambda,H_j}^D P_{H_j} \langle n_{\Lambda,H_j} \rangle] / \langle n_\Lambda^{(total)} \rangle. \quad (1)$$

where $\langle n_\Lambda^{(1)} \rangle$ and $\langle n_\Lambda^{(2)} \rangle$ are the average numbers of the Λ 's of the origins (1) and (2) mentioned above respectively, and $\langle n_{\Lambda,H_j} \rangle$ is that from the decay of H_j 's; P_{H_j} is the polarization of H_j ; $\langle n_\Lambda^{(total)} \rangle$ is the total number of Λ 's. These average numbers are determined by the hadronization mechanisms, and we just use the results obtained from LEPTO. $t_{\Lambda,u}^F$ is the fragmentation polarization transfer factor, which is the probability for the polarization of u -quark to be transferred to Λ if the Λ contains this u -quark. It is different in the SU(6) or the DIS picture. t_{H_i,H_j}^D is the decay polarization transfer factor determined by the decay process. For details of $t_{\Lambda,u}^F$ and t_{H_i,H_j}^D , see e.g. [12] and the references given there.

After the calculations, we obtain the longitudinal polarization of Λ in $\nu_\mu p \rightarrow \mu^- \Lambda X$ and $\nu_\mu n \rightarrow \mu^- \Lambda X$ at $E_\nu = 44\text{GeV}$ as shown in Figs.2a and 2b. To show the influence of the nucleon remnant fragmentation, in the figures, we show also the results in the case when only the struck quark fragmentation is taken into account. From Fig.2a, we see

that the contributions from the (uu) -diquark fragmentation are indeed very large, even at $x_F > 0$. In fact, the characteristic features of the results in the region around $x_F \sim 0$ are mainly determined by the (uu) -diquark fragmentation. E.g., we see that, in this region, $P_{\Lambda(L)}$ is negative and $|P_{\Lambda(L)}|$ increases with decreasing x_F . This is determined by the decay contribution of the Σ^{*+} 's which contain the (uu) diquarks. It contributes negatively to $P_{\Lambda(L)}$ and the contribution increases with the decreasing x_F . Compare the results in Fig.2a with those in Fig.2b, we see that, $P_{\Lambda(L)}$ for $\nu_\mu n \rightarrow \mu^- \Lambda X$ differs greatly from that for $\nu_\mu p \rightarrow \mu^- \Lambda X$. It is very small in the whole x_F region. This is because the contributions from the decay of the heavier hyperons which contain the $(ud)_1$ or a u - or a d -quark from $(ud)_1$ are different in sign and cancel with each other. Since NOMAD has approximately an isobar target, the NOMAD data¹⁴ corresponds approximately to a mixture of (2/3) n- and (1/3) p-targets. We add the results with these weights and compare the obtained results with the data¹⁴ in Fig.2c. We see that the qualitative features of the results agree with the data¹⁴.

We also made similar calculations for $P_{\Sigma^+(L)}$ in $\nu_\mu N \rightarrow \mu^- \Sigma^+ X$ and obtained the results in Fig.3. The advantage to study Σ^+ instead of Λ is that the decay contributions to Σ^+ are much smaller. The major contribution to $\nu_\mu p \rightarrow \mu^- \Sigma^+ X$ is from the (uu) fragmentation. The Σ^+ 's which contain the (uu) -diquark dominate for most x_F . This is why we see that the obtained $P_{\Sigma^+(L)}$ is negative and its absolute value is very large for $\nu_\mu p \rightarrow \mu^- \Sigma^+ X$. These can be checked by future experiments.

It should be noted that, almost at the same time, another group has carried out¹⁸ the similar calculations as presented in this paper independently using a different model for the spin transfer in the fragmentation of diquarks. The aim of the authors was to use Λ polarization in such process to study the polarization of the strange sea in nucleon. Assuming that there is a correlation between the polarization of the strange sea and that of the struck quark (which was not included in our calculations), they obtained a good fit to the NOMAD data¹⁴. The authors also found similar effect which confirms the conclusion that we present above and briefly reported earlier in the conference talks¹⁰. We also note that, in our

calculations, we used the default set of parameters in JETSET, and such set of parameters may not, as pointed out in [18], reproduce the data on the average numbers of the strange baryons produced in the process. However, since we are interested in the relative contributions of Λ from the different sources (1) to (4) mentioned above, this may be not very sensitive to our results in particular the qualitative conclusions we reached above. More precisely, there should be little influence from the parameters such as the overall strange suppression etc. But there can be some influence from the parameters such as decuplet baryon production suppression etc since a large part of the contribution from the diquark fragmentation come from Σ^* decay. To see whether this is true, we tuned the related parameters and repeated the calculations. We found out that there are indeed some influence on the quantitative results but the qualitative features remain.

In summary, we show that the products from the fragmentation of the struck quark and those of the nucleon remnant can not be separated from each other in the NOMAD energy region. The latter gives the dominant contribution in the region around $x_F \sim 0$ and the characteristic features of the hyperon polarization in this region are determined by this contribution. We made a rough estimation of the Λ polarizations by taking the fragmentation of the nucleon remnant into account and the results are consistent with the NOMAD data. Further predictions for $\nu_\mu N \rightarrow \mu^- \Sigma^+ X$ are made.

We are in debt to Dr. D. Naumov, who suggested us to make the research in the target fragmentation region and continuously communicated with us on different aspects. This work was supported in part by the National Science Foundation of China (NSFC), the Education Ministry of China under Huo Ying-dong Foundation, and the Postdoctoral Science Foundation of China.

REFERENCES

1. R.L. Jaffe, and X. Ji, Phys. Rev. Lett. **67** , 552 (1991); Nucl. Phys. B**375**, 527 (1992); M. Burkardt and R.L. Jaffe, Phys. Rev. Lett. **70**, 2537 (1993); R.L. Jaffe, Phys. Rev. D**54**, R6581 (1996).
2. G.Gustafson, J.Häkkinen, Phys.Lett.B**303**,350(1993).
3. C. Boros, and Z. Liang, Phys. Rev. D**57**, 4491 (1998).
4. A. Kotzinian, A. Bravar, D. von Harrach, Eur. Phys. J. C**2**, 329 (1998).
5. D. Ashery, H.J. Lipkin, Phys. Lett. B**469**, 263 (1999); hep-ph/0002144.
6. B.Q. Ma, I. Schmidt, and J.J. Yang, Phys. Rev. D**61**, 034017 (2000).
7. M.Anselmino, M.Boglione and F.Murgia, Phys. Lett. B**481**, 253 (2000).
8. C. Liu and Z. Liang, Phys. Rev. D**62**, 094001 (2000).
9. B.Q. Ma, I. Schmidt, J. Soffer, and J.J. Yang, Phys. Rev. D**62**, 114009 (2000).
10. C. Liu, Q. Xu, and Z. Liang, Phys. Rev. D**64**, 073004 (2001); Z. Liang, in Proc. of the 31st Inter. Sympo. on Multiparticle Dynamics, Datong, 2001, World Scientific (2002), p.78, hep-ph/0111403; and Proc. 3rd Circum-Pan-Pacific Sympo. on High Energy Spin Phys., Beijing, 2001, hep-ph/0205017; C. Liu, Q. Xu, Z. Liang, *ibid.*
11. M.Anselmino, M.Boglione, U.D'Alesio, and F.Murgia, Eur. Phys. J. C**21**, 501 (2001).
12. Xu Qing-hua, Liu Chun-xiu and Liang Zuo-tang, Phys. Rev. D**65**, 114008 (2002).
13. HERMES Collaboration, A. Airapetian *et al.*, Phys. Rev. D**64**, 112005 (2001).
14. NOMAD Collaboration, P. Astier *et al.*, Nucl. Phys. B**588**, 3 (2000); **605**, 3 (2001); D. Naumov, in Proc. of the 14th Inter. Sympo. on Spin Phys., AIP Conf. Proc. 570 (2001), p.489, hep-ph/0101325.
15. M.L. Mangano *et al.*, hep-ph/0105155.

16. G Ingelman, A.Edin, J.Rathsman, Comp. Phys. Comm. **101**, 108 (1997).
17. B. Anderson, *et al.*, Phys. Rep. **97**, 31 (1983).
18. J.Ellis, A. Kotznian, and D. Naumov, hep-ph/0204206.

FIGURES




Fig. 1. Different contributions to Λ in $\nu_\mu p \rightarrow \mu^- \Lambda X$ at $E_\nu = 44\text{GeV}$. Here, the solid, dotted, dashed and dash-dotted lines denote respectively (1) directly produced and contain the struck quark; (2) directly produced and contain a u -quark in $(uu)_1$; (3) decay products of hyperons which contain the struck quark; (4) decay products of hyperons which contain the $(uu)_1$ or a u in the $(uu)_1$.




Fig. 2. $P_{\Lambda(L)}$ as a function of x_F in $\nu_\mu N \rightarrow \mu^- \Lambda X$ at $E_\nu = 44\text{ GeV}$ when both the contributions from the fragmentation of the struck quark and that of the nucleon remnant are taken into account (solid and dotted lines), compared to the results when only the struck quark fragmentation is taken into account (dashed and dash-dotted lines). The data are taken from [14].




Fig. 3. $P_{\Sigma^+(L)}$ as a function of x_F in $\nu_\mu N \rightarrow \mu^- \Sigma^+ X$ at $E_\nu = 44$ GeV when both the contributions from the fragmentation of the struck quark and that of the nucleon remnant are taken into account (solid and dotted lines), compared to the results when only the struck quark fragmentation is taken into account (dashed and dash-dotted lines).